

d his

(FILE 'HOME' ENTERED AT 09:35:52 ON 04 MAR 2005)

FILE 'STNGUIDE' ENTERED AT 09:35:58 ON 04 MAR 2005

FILE 'REGISTRY' ENTERED AT 09:36:11 ON 04 MAR 2005

L1           1 S NOOTKATONE/CN  
L2           3 S NOOTKATONE/CN OR CREDENE OR ZIZANOL/CN OR BICYCLOVETIVENOL/CN  
L3           31 S NOOTKATONE/CN OR CEDRENE OR ZIZANOL/CN OR BICYCLOVETIVENOL/CN

FILE 'CAPPLUS, USPATFULL' ENTERED AT 09:37:33 ON 04 MAR 2005

L4           1863 S NOOTKATONE OR CEDRENE OR ZIZANOL OR BICYCLOVETIVENOL  
L5           4310 S (CONTROL? OR KILL? OR REPEL?) (2A) (TICK? OR COCKROACH? OR CO  
L6           7 S L4 (P) L5  
L7           1 S TOPICAL? (3A) L4  
L8           68196 S (APPL? OR TREAT? OR CONTACT?) (2A) (SKIN OR EPIDERM? OR DERMA  
L9           2 S L8 (P) L4

FILE 'REGISTRY' ENTERED AT 10:10:15 ON 04 MAR 2005

L10          1 S ZIZANOL/CN  
L11          1 S BICYCLOVETIVENOL/CN

FILE 'CAPPLUS, USPATFULL' ENTERED AT 10:10:54 ON 04 MAR 2005

L12          23 S ZIZANOL OR L10  
L13          21 S BICYCLOVETIVENOL OR L11  
L14          9 S L12 AND L13  
L15          7 S (COMPOSITION OR FORMULATION) (3A) L14  
L16          2 S L14 NOT L15  
L17          14 S L12 NOT L14  
L18          14 S L12 NOT L14  
L19          12 S L13 NOT L14  
L20          12 S L19 NOT L17

=>

L15 ANSWER 3 OF 7 CAPIUS COPYRIGHT 2005 ACS on STN  
ACCESSION NUMBER: 1989:44730 CAPIUS  
DOCUMENT NUMBER: 110:44730  
TITLE: GC-MS and GC-MS-MS analysis of a complex essential oil  
AUTHOR(S): Cazaussus, Annie; Pes, R.; Sellier, Nicole; Tabet, J. C.  
CORPORATE SOURCE: Lab. Spectrometrie Masse, CERCOA, Paris, F-75231/05, Fr.  
SOURCE: Chromatographia (1988), 25(10), 865-9  
CODEN: CHRGB7; ISSN: 0009-5893  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB The alc. portion of a vetiver oil, the gas chromatogram (GC) of which shows more than 100 components, was analyzed by combined GC-mass spectrometry (MS) with different ionization methods. The specificity of the method was improved by using combined GC-tandem MS in the collision-activated-dissociation mode. The vapor components were identified as  $\beta$ -vetivenene, khusimone, zizanal, **zizanol**, **bicyclovetivenol**,  $\beta$ -vetivone, and  $\alpha$ -vetivone.

AB The alc. portion of a vetiver oil, the gas chromatogram (GC) of which shows more than 100 components, was analyzed by combined GC-mass spectrometry (MS) with different ionization methods. The specificity of the method was improved by using combined GC-tandem MS in the collision-activated-dissociation mode. The vapor components were identified as  $\beta$ -vetivenene, khusimone, zizanal, **zizanol**, **bicyclovetivenol**,  $\beta$ -vetivone, and  $\alpha$ -vetivone.

ST vetiver oil **compn**; sesquiterpene vetiver oil

IT Oils, essential

RL: PRP (Properties)  
(**compn.** of, gas chromatog.-mass spectrometry of)

IT Oils, essential

RL: PRP (Properties)  
(vetiver, **compn.** of, gas chromatog.-mass spectrometry of)

IT 5957-31-3, **Bicyclovetivenol** 15764-04-2,  
 $\alpha$ -Vetivone 18444-79-6,  $\beta$ -Vetivone 27840-40-0,  
 $\beta$ -Vetivenene 28102-79-6 30557-76-7, Khusimone  
82509-29-3, Zizanal

RL: BIOL (Biological study)  
(of vetiver oil, gas chromatog.-mass spectrometry of)

L6 ANSWER 3 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1997:447331 CAPLUS

DOCUMENT NUMBER: 127:77365

TITLE: Acaricides containing phenyl isothiocyanate

INVENTOR(S): Ishida, Hirohiko; Izumi, Tasuku

PATENT ASSIGNEE(S): Kao Corp., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 7 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 09157116	A2	19970617	JP 1995-346872	19951212
			JP 1995-346872	19951212

PRIORITY APPLN. INFO.:

AB The acaricides contain Ph isothiocyanate (I) and optionally perfumes selected from (A) sesquiterpenes, (B) essential oils containing sesquiterpenes, (C) alicyclic compds., and (D) aromatic compds. as pungent odor-masking components. I was mixed with a perfume composition containing **cedrene** 40, cedarwood oil 10, vetiverol 5, caryophyllene 5, and hinoki oil 35 parts at a weight ratio of 1:19 to give an acaricide, which was placed in a container (at 250 mg/50 g culture media) to show 100% control of **ticks** after 4 wk at 22-25° and relative humidity 75-85%. A flooring material was sprayed with the acaricide and left at room temperature for 3 days to show no pungent odor. Formulation examples are given.

AB The acaricides contain Ph isothiocyanate (I) and optionally perfumes selected from (A) sesquiterpenes, (B) essential oils containing sesquiterpenes, (C) alicyclic compds., and (D) aromatic compds. as pungent odor-masking components. I was mixed with a perfume composition containing **cedrene** 40, cedarwood oil 10, vetiverol 5, caryophyllene 5, and hinoki oil 35 parts at a weight ratio of 1:19 to give an acaricide, which was placed in a container (at 250 mg/50 g culture media) to show 100% control of **ticks** after 4 wk at 22-25° and relative humidity 75-85%. A flooring material was sprayed with the acaricide and left at room temperature for 3 days to show no pungent odor. Formulation examples are given.

L6 ANSWER 4 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN  
ACCESSION NUMBER: 1996:400558 CAPLUS  
DOCUMENT NUMBER: 125:51527  
TITLE: Cockroach repellents containing sesquiterpenes  
INVENTOR(S): Harima, Shoichi; Komai, Koichiro; Myake, Takayasu  
PATENT ASSIGNEE(S): Tokiwa Kanpo Pharma, Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 9 pp.  
CODEN: JKXXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	-----	-----	-----	-----
JP 08081306	A2	19960326	JP 1994-217066	19940912
PRIORITY APPLN. INFO.:			JP 1994-217066	19940912

AB Cockroach repellents contain (A) sesquiterpenes containing  $\geq 1$  groups chosen from CMe2OH, CHMeCH2OH, CH2OH, and cyclic ketones and  $\geq 1$  double bonds or (B) sesquiterpene alcs. containing 1 group chosen from CMe2OH and CHMeCH2OH and terminal methylene as active ingredients. Elemol at 1.0 g/m<sup>2</sup> showed 71-100% repellency against Blattella germanica and Periplaneta fuliginosa at 24 h after the application.

IT 473-15-4P,  $\beta$ -Eudesmol 639-99-6P, Elemol 4674-50-4P,  
**Nootkatone** 15051-81-7P, 10-epi- $\gamma$ -Eudesmol 16223-63-5P,  
Khusimol 18444-79-6P,  $\beta$ -Vetivone 28102-68-3P, Vetiselinol  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BUU (Biological use, unclassified); PUR (Purification or recovery); BIOL (Biological study); PREP (Preparation);  
USES (Uses)

(cockroach repellents containing sesquiterpenes)